

# 1 - 4 Weeks Questions



Patient Name:	DOB:	Date:
---------------	------	-------

**Any questions or concerns today? Please circle all that apply.**

- |   |   |   |
|---|---|---|
| Eating  | Y | N |
| Elimination (voiding/stooling)  | Y | N |
| Sleeping  | Y | N |
| Developmental   | Y | N |
| Hearing/Vision  | Y | N |
| Skin  | Y | N |
| Vaccines  | Y | N |
| Other   | Y | N |
| <b>Has your baby been fussy?</b>  | Y | N |
| <b>Can you tell when your baby needs to eat?</b>                          | Y | N |
| <b>Can you tell when your baby needs to sleep?</b>                        | Y | N |
| <b>Do you know what shaken baby syndrome is?</b>                          | Y | N |
| <b>Do you have pets?</b>  | Y | N |
| Cat   | Y | N |
| Dog   | Y | N |
| Other   | Y | N |
| <b>Does your baby have any exposure to smoke?</b>                         | Y | N |
| <b>Do you have a rear facing car seat?</b>                                | Y | N |
| <b>Are you returning to work/school?</b>                                  | Y | N |
| <b>Do you have working smoke detectors and carbon monoxide detectors?</b> | Y | N |
| <b>Do you have any guns in the home?</b>                                  | Y | N |
| Are the guns unloaded?  | Y | N |
| Are the guns locked?  | Y | N |
| Is ammunition stored separately?  | Y | N |

**SEE REVERSE FOR ADDITIONAL QUESTIONS**

# 1 - 4 Weeks Questions



Patient Name:

DOB:

Date:

## Food Insecurity & Transportation Questions (mark your answer):

- Within the past 12 months, you worried that your food would run out before you got money to buy more.
  - Often true
  - Sometimes true
  - Never true
- Within the past 12 months, the food you bought just didn't last and you didn't have money to get more.
  - Often true
  - Sometimes true
  - Never true
- In the past 12 months, has lack of transportation kept you from medical appointments, meetings, working or from getting things needed for daily living (mark all that apply)?
  - Yes, it has kept me from medical appointments or getting medications
  - Yes, it has kept me from non-medical meetings, appointments, work or getting things that I need
  - No

# Edinburgh Postnatal Depression Scale<sup>1</sup> (EPDS)

Name: \_\_\_\_\_

Address: \_\_\_\_\_

Your Date of Birth: \_\_\_\_\_

\_\_\_\_\_

Baby's Date of Birth: \_\_\_\_\_

Phone: \_\_\_\_\_

---

As you are pregnant or have recently had a baby, we would like to know how you are feeling. Please check the answer that comes closest to how you have felt **IN THE PAST 7 DAYS**, not just how you feel today.

Here is an example, already completed.

I have felt happy:

- Yes, all the time
- Yes, most of the time      This would mean: "I have felt happy most of the time" during the past week.
- No, not very often      Please complete the other questions in the same way.
- No, not at all

In the past 7 days:

- |   |   |
|---|---|
| 1. I have been able to laugh and see the funny side of things | *6. Things have been getting on top of me   |
| <input type="checkbox"/> As much as I always could            | <input type="checkbox"/> Yes, most of the time I haven't been able to cope at all |
| <input type="checkbox"/> Not quite so much now                | <input type="checkbox"/> Yes, sometimes I haven't been coping as well as usual    |
| <input type="checkbox"/> Definitely not so much now           | <input type="checkbox"/> No, most of the time I have coped quite well             |
| <input type="checkbox"/> Not at all                           | <input type="checkbox"/> No, I have been coping as well as ever                   |
| 2. I have looked forward with enjoyment to things             | *7. I have been so unhappy that I have had difficulty sleeping                    |
| <input type="checkbox"/> As much as I ever did                | <input type="checkbox"/> Yes, most of the time                                    |
| <input type="checkbox"/> Rather less than I used to           | <input type="checkbox"/> Yes, sometimes   |
| <input type="checkbox"/> Definitely less than I used to       | <input type="checkbox"/> Not very often   |
| <input type="checkbox"/> Hardly at all                        | <input type="checkbox"/> No, not at all   |
| *3. I have blamed myself unnecessarily when things went wrong | *8. I have felt sad or miserable  |
| <input type="checkbox"/> Yes, most of the time                | <input type="checkbox"/> Yes, most of the time                                    |
| <input type="checkbox"/> Yes, some of the time                | <input type="checkbox"/> Yes, quite often   |
| <input type="checkbox"/> Not very often                       | <input type="checkbox"/> Not very often   |
| <input type="checkbox"/> No, never                            | <input type="checkbox"/> No, not at all   |
| 4. I have been anxious or worried for no good reason          | *9. I have been so unhappy that I have been crying                                |
| <input type="checkbox"/> No, not at all                       | <input type="checkbox"/> Yes, most of the time                                    |
| <input type="checkbox"/> Hardly ever                          | <input type="checkbox"/> Yes, quite often   |
| <input type="checkbox"/> Yes, sometimes                       | <input type="checkbox"/> Only occasionally  |
| <input type="checkbox"/> Yes, very often                      | <input type="checkbox"/> No, never  |
| *5. I have felt scared or panicky for no very good reason     | *10. The thought of harming myself has occurred to me                             |
| <input type="checkbox"/> Yes, quite a lot                     | <input type="checkbox"/> Yes, quite often   |
| <input type="checkbox"/> Yes, sometimes                       | <input type="checkbox"/> Sometimes  |
| <input type="checkbox"/> No, not much                         | <input type="checkbox"/> Hardly ever  |
| <input type="checkbox"/> No, not at all                       | <input type="checkbox"/> Never  |

Administered/Reviewed by \_\_\_\_\_ Date \_\_\_\_\_

<sup>1</sup>Source: Cox, J.L., Holden, J.M., and Sagovsky, R. 1987. Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* 150:782-786 .

<sup>2</sup>Source: K. L. Wisner, B. L. Parry, C. M. Piontek, Postpartum Depression N Engl J Med vol. 347, No 3, July 18, 2002, 194-199

Users may reproduce the scale without further permission providing they respect copyright by quoting the names of the authors, the title and the source of the paper in all reproduced copies.





# BRIGHT FUTURES HANDOUT ► PARENT

## FIRST WEEK VISIT (3 TO 5 DAYS)

Here are some suggestions from Bright Futures experts that may be of value to your family.

### ✓ HOW YOUR FAMILY IS DOING

- If you are worried about your living or food situation, talk with us. Community agencies and programs such as WIC and SNAP can also provide information and assistance.
- Tobacco-free spaces keep children healthy. Don't smoke or use e-cigarettes. Keep your home and car smoke-free.
- Take help from family and friends.

### ✓ FEEDING YOUR BABY

- Feed your baby only breast milk or iron-fortified formula until he is about 6 months old.
- Feed your baby when he is hungry. Look for him to
  - Put his hand to his mouth.
  - Suck or root.
  - Fuss.
- Stop feeding when you see your baby is full. You can tell when he
  - Turns away
  - Closes his mouth
  - Relaxes his arms and hands
- Know that your baby is getting enough to eat if he has more than 5 wet diapers and at least 3 soft stools per day and is gaining weight appropriately.
- Hold your baby so you can look at each other while you feed him.
- Always hold the bottle. Never prop it.

#### If Breastfeeding

- Feed your baby on demand. Expect at least 8 to 12 feedings per day.
- A lactation consultant can give you information and support on how to breastfeed your baby and make you more comfortable.
- Begin giving your baby vitamin D drops (400 IU a day).
- Continue your prenatal vitamin with iron.
- Eat a healthy diet; avoid fish high in mercury.

#### If Formula Feeding

- Offer your baby 2 oz of formula every 2 to 3 hours. If he is still hungry, offer him more.

### ✓ HOW YOU ARE FEELING

- Try to sleep or rest when your baby sleeps.
- Spend time with your other children.
- Keep up routines to help your family adjust to the new baby.

### ✓ BABY CARE

- Sing, talk, and read to your baby; avoid TV and digital media.
- Help your baby wake for feeding by patting her, changing her diaper, and undressing her.
- Calm your baby by stroking her head or gently rocking her.
- *Never hit or shake your baby.*
- Take your baby's temperature with a rectal thermometer, not by ear or skin; a fever is a rectal temperature of 100.4°F/38.0°C or higher. Call us anytime if you have questions or concerns.
- Plan for emergencies: have a first aid kit, take first aid and infant CPR classes, and make a list of phone numbers.
- Wash your hands often.
- Avoid crowds and keep others from touching your baby without clean hands.
- Avoid sun exposure.

**Helpful Resources:** Smoking Quit Line: 800-784-8669 | Poison Help Line: 800-222-1222

Information About Car Safety Seats: [www.safercar.gov/parents](http://www.safercar.gov/parents) | Toll-free Auto Safety Hotline: 888-327-4236

# FIRST WEEK VISIT (3 TO 5 DAYS)—PARENT



## SAFETY

- Use a rear-facing-only car safety seat in the back seat of all vehicles.
- Make sure your baby always stays in his car safety seat during travel. If he becomes fussy or needs to feed, stop the vehicle and take him out of his seat.
- Your baby's safety depends on you. Always wear your lap and shoulder seat belt. Never drive after drinking alcohol or using drugs. Never text or use a cell phone while driving.
- Never leave your baby in the car alone. Start habits that prevent you from ever forgetting your baby in the car, such as putting your cell phone in the back seat.
- Always put your baby to sleep on his back in his own crib, not your bed.
  - Your baby should sleep in your room until he is at least 6 months old.
  - Make sure your baby's crib or sleep surface meets the most recent safety guidelines.
- If you choose to use a mesh playpen, get one made after February 28, 2013.
- Swaddling is not safe for sleeping. It may be used to calm your baby when he is awake.
- Prevent scalds or burns. Don't drink hot liquids while holding your baby.
- Prevent tap water burns. Set the water heater so the temperature at the faucet is at or below 120°F /49°C.

## WHAT TO EXPECT AT YOUR BABY'S 1 MONTH VISIT

### We will talk about

- Taking care of your baby, your family, and yourself
- Promoting your health and recovery
- Feeding your baby and watching her grow
- Caring for and protecting your baby
- Keeping your baby safe at home and in the car

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents, 4th Edition*

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.